

선광일 (Kwang-Il Seon) UST / KASI

For an isotropic, and extremely relativistic electron distribution, we can use energy instead of lacksquaremomentum:

$$E = \sqrt{(pc)^{2} + (mc^{2})^{2}} \approx pc \implies d^{3}p = 4\pi p^{2}dp = \frac{4\pi}{c^{3}}E^{2}dE$$

$$f(p)4\pi p^{2}dp = N(E)dE \implies f(p) = \frac{N(E)dE}{4\pi p^{2}dp} = \frac{N(E)}{(4\pi/c^{3})E^{2}} \implies \begin{cases} d^{3}pf(p) = dEE^{2}\frac{N(E)}{E^{2}} \\ d^{3}pf(p^{*}) = dEE^{2}\frac{N(E^{*})}{E^{*2}} \\ d^{3}pf(p^{*}) = dEE^{2}\frac{N(E^{*})}{E^{*2}} \end{cases}$$
Then,
$$\alpha_{\nu} = \frac{c^{2}}{8\pi h\nu^{3}}\int dEP(\nu, E)E^{2}\left[\frac{N(E-h\nu)}{(E-h\nu)^{2}} - \frac{N(E)}{E^{2}}\right] \longleftarrow \begin{bmatrix} \alpha_{\nu} = \frac{c^{2}}{8\pi h\nu^{3}}\int d^{3}p_{2}\left[f(p_{2}^{*}) - f(p_{2})\right]P(\nu, E_{2}) \end{cases}$$

Assume that  $h\nu \ll E$  (in fact, a necessary condition for the application of classical electrodynamics) and expand to first order in  $h\nu$ . Taylor expansion:

$$\alpha_{\nu} = -\frac{c^2}{8\pi\nu^2} \int dEP(\nu, E) E^2 \frac{\partial}{\partial E} \left[ \frac{N(E)}{E^2} \right] \qquad \qquad \frac{N(E - h\nu)}{(E - h\nu)^2} \approx \frac{N(E)}{E^2} - h\nu \frac{\partial}{\partial E} \left[ \frac{N(E)}{E^2} \right] + \mathcal{O}\left((h\nu)^2\right)$$

For a power law distribution of particles: 

• For a power law distribution of particles:  

$$N(E) = CE^{-p}$$

$$-E^{2}\frac{d}{dE}\left[\frac{N(E)}{E^{2}}\right] = (p+2)CE^{-(p+1)}$$

$$= \frac{(p+2)N(E)}{E}$$

$$\alpha_{\nu} = \frac{(p+2)c^{2}}{8\pi\nu^{2}}\int dEP(\nu, E)\frac{N(E)}{E}$$

$$\propto \nu^{-2}\int dEF(x)\frac{E^{-p}}{E} \quad \leftarrow \text{ set } x = \frac{\omega}{\omega_{c}} \propto \nu\gamma^{-2} \propto \nu E^{-2}$$

$$\propto \nu^{-2}\int \nu^{1/2}x^{-3/2}dxF(x)\nu^{-(p+1)/2}x^{(p+1)/2}$$

$$\alpha_{\nu} \propto B^{(p+2)/2}\nu^{-(p+4)/2}$$

# Note $\alpha_{\nu} \propto \nu^{-(p+4)/2}$ indicates that the synchrotron emission is optically thick at low frequencies and optically thin at high frequencies.

The source function is

$$S_{\nu} = \frac{j_{\nu}}{\alpha_{\nu}} = \frac{P(\nu)}{4\pi\alpha_{\nu}} \propto B^{-1/2}\nu^{5/2} \qquad \qquad P(\nu) \propto B^{(p+1)/2}\nu^{-(p-1)/2} \\ \alpha_{\nu} \propto B^{(p+2)/2}\nu^{-(p+4)/2} \end{cases}$$
For optically thin synchrotron emission, 
$$I_{\nu} = \int j_{\nu} ds \propto B^{(p+1)/2}\nu^{-(p-1)/2} \\ I_{\nu} = S_{\nu} \propto B^{-1/2}\nu^{5/2} \qquad \text{high frequency}$$

Therefore, the synchrotron spectrum from a power-law distribution of electrons has a shape like the following figure.



- Observations of the self-absorption part could determine *B*.
- Observations of the thin part can then determine the proportional constant *K* and the electron slope *p*.

## Polarization of Synchrotron Radiation



Electrons rotate counterclockwise when viewed from the positive tip of the **B** axis.

gyrofrequency : 
$$\omega_{\rm B} = \frac{eB}{\gamma m_e c}$$
  
gyroradius :  $r_{\rm B} = \frac{v_{\perp}}{\omega_{\rm B}} (v_{\perp} = v \sin \alpha)$ 



[left] Non-relativistic cyclotron motion. When viewed in orbital plane, radiation is 100% linearly polarized with electric vector oscillating perpendicular to magnetic field **B**. Viewed from along **B**, emission is 100% circularly polarized. Note that the electric vector depends only on the observer's direction and is independent of the pitch angle.

[right] For **relativistic motion**, radiation is beamed into direction of motion. The emission for a single electron is effectively confined to within a small angle  $1/\gamma$  of **v** (The electric field depends both on the observer's direction and pitch angle). The fourth Stokes parameter is an odd function of the angle between **n** and **v**. The number of electrons passing with an pitch angle  $\alpha$  is the same as that with  $-\alpha$ . These two components of circular polarization effectively cancel almost, whereas linear polarization largely survives.

### **Cooling Time Scale**

• Emission as synchrotron radiation of relativistic electrons, characteristic frequency:

$$\nu_c \simeq \frac{1}{2\pi} \gamma^2 \frac{eB}{m_e c} \simeq 280 \left(\frac{B}{10^{-4}\,\mathrm{G}}\right) \gamma^2\,\mathrm{Hz}$$

for average  $B \sim 10^{-4}$  G in Crab Nebula

#### • Optical Emission:

Optical emission (HST) at  $\nu \sim 5 \times 10^{14}$  Hz requires electrons with  $\gamma \sim 10^6$ . Cooling time scale  $t_{cool} \sim 2500 (10^6/\gamma)$  yr  $\gtrsim t_{age}$  age of Nebula.

### • X-ray Emission:

Chandra (ACIS, 0.2-10 keV) X-ray emission  $\nu \sim 10^{17}$  Hz requires  $\gamma \sim 10^7$ , electrons cool quicker by a factor ~ 10.

X-ray emission spatially less extended.

 $t_{\rm cool} < t_{\rm age} \sim 950$  yr of Nebula, need continuous supply of fresh electrons.

#### • Radio Emission:

Crab Nebula also bright in radio (NRAO,  $\nu \sim 5 \times 10^9$  Hz), less energetic electrons needed,  $\gamma \sim 5 \times 10^3$ , size constrained by the age of Nebula

### Astrophysical Examples: Synchrotron Emission from Crab Nebula

6



Crab Nebula ( $d \sim 2$  kpc) caused by SN explosion in 1054 A.D. - composite image. Chandra X-ray [blue], HST optical [red and yellow], Spitzer infrared [purple]. X-ray image is smaller than others as extremely energetic electrons emitting X-rays radiate away their energy more quickly than lower-energy electrons emitting in optical and infrared [Credits: NASA]

• Crab nebula - SED

7

Dots: modified blackbody with T = 46 K. Thin dashed line: emission at mm wavelengths Thick dashed line: synchrotron emission The solid lines are a fit to the data for a synchrotron self-Compton (SSC) model for a nebular magnetic field of 16 nT (= 0.16 mG) with contributions from a millimeter synchrotron (radio) region and from nebular dust.

The peak at TeV energies is attributed to synchrotron photons that have been upscattered via inverse Compton scattering.



• Crab nebula - polarization



Photographs of the Crab nebula in polarized light with the polarizer at different orientations. The arrows show the directions or planes of the transmitted transverse electric vector. Note the changing brightness pattern from photo to photo. The nebula has angular size  $4' \times 6'$  and is ~6000 LY distant from the solar system. North is up and east to the left. The pulsar is the southwest (lower right) partner of the doublet at the center of the nebula. [Palomar Observatory/CalTech]

### Astrophysical Examples: Spinning neutron star

• Spinning neutron star



Spinning neutron star with offset magnetic pole. Electrons spiral outward along magnetic field lines and beam radiation in the polar direction. (The spiraling electrons illustrated would not radiate into the beam.)

## Astrophysical Examples: Galactic radio emission

• Galactic radio synchrotron radiation at 150 MHz.

The diffuse radio emission observed over the whole sky is the result of the radiation by **cosmic-ray** electrons interacting with interstellar magnetic fields. It is concentrated toward the galactic plane, exhibits a power-law character, and has the expected polarization.



The contours are brightness temperatures (K). Most of this radiation is synchrotron radiation from electrons spiraling around magnetic fields in interstellar space within the Galaxy [Landecker & Wielebinski, 1970]

# Compton Scattering

# **Thomson & Compton Scattering**

- The simplest interaction between photons and free electrons is scattering.
  - **Thomson scattering:** When the energy of the incoming photons (as seen in the coming frame of the electron) is small with respect to the electron rest mass-energy, the process is called Thomson scattering.

$$\begin{aligned} \epsilon &= \epsilon_1 \\ \frac{d\sigma_T(\Omega)}{d\Omega} &= \frac{1}{2}r_0^2 \left(1 + \cos^2 \theta\right) \\ \sigma_T &= \frac{8\pi}{3}r_0^2 \end{aligned} \qquad \begin{aligned} \epsilon &= \text{energy of the incident photon} \\ \epsilon_1 &= \text{energy of the scattered photon} \\ r_0 &= \frac{e^2}{m_e c^2} \\ \end{aligned} \qquad \begin{aligned} Thomson \text{ scattering condition in the rest frame:} \\ \epsilon' &\ll m_e c^2 = 0.5 \text{MeV} \end{aligned} \end{aligned}$$

- When  $\epsilon = \epsilon_1$ , the scattering is called **coherent or elastic**.
- **Compton scattering:** As the energy of the incoming photons is comparable or greater than the electron rest mass-energy, it is called Compton scattering and a quantum treatment is necessary (Klein-Nishina regime).

# [Compton Scattering: Scattering from Electrons at Rest]

#### • <u>Compton scattering:</u>

However, a photon carries momentum  $h\nu/c$  and energy  $h\nu$ .

Quantum effects appear in two ways.

(1) The scattering will no longer be elastic ( $\epsilon_1 \neq \epsilon_2$ ) because of the recoil of the electron.

(2) The cross sections are altered by the quantum effects.

• Conservation of momentum and energy (for the case in which **the electron is initially at rest**)

Let the initial and final four-momenta of the photon:

$$\vec{P}_{\gamma i} = (\epsilon/c)(1, \mathbf{n}_i), \ \vec{P}_{\gamma f} = (\epsilon_1/c)(1, \mathbf{n}_f)$$

The initial and final momenta of the electron are:

$$\vec{P}_{ei} = (mc, \mathbf{0}), \ \vec{P}_{ef} = (E/c, \mathbf{p})$$

Then, the conservation of momentum and energy is expressed by

$$\vec{P}_{ei} + \vec{P}_{\gamma i} = \vec{P}_{ef} + \vec{P}_{\gamma f}$$

Here, e and  $\gamma$  denote the electron and photon, respectively. i and f represent the initial and final states.



• Rearranging terms and squaring gives  $\left|\vec{P}_{ef}\right|^2 = \left|\vec{P}_{ei} + \vec{P}_{\gamma i} - \vec{P}_{\gamma f}\right|^2$ 

There is a wavelength change of the order of  $\lambda_c$  upon scattering. For long wavelengths  $\lambda \gg \lambda_c$  (i.e.,  $h\nu \ll mc^2$ ), the scattering is closely elastic.



Scattered photons energies are shown as a function of the scattering angle, for different incoming photon energies.

Note that, for  $x \gg 1$  and for large scattering angle, the scattered photon energies becomes  $x_1 \sim 1/2$ , independent of the initial photon energy *x*.



Compton scattering becomes less efficient at high energies. Approximations:

- nonrelativisitic regime:
- extreme relativisitic regime:

$$\sigma \approx \sigma_T \left( 1 - 2x + \frac{26x^2}{5} + \cdots \right), \quad x \ll 1$$
$$\sigma \approx \frac{3}{8} \sigma_T \frac{1}{x} \left( \ln 2x + \frac{1}{2} \right), \quad x \gg 1$$

 $(m_e c^2 = 511 \text{ keV})$ 

## [Inverse Compton Scattering: Scattering from Electrons in Motion]

 $\mathbf{n}' = d\mathbf{i}$ 

 $\mathbf{n}_1' = \mathbf{d}$ 

- **Inverse Compton Scattering**: Whenever the moving electron has sufficient kinetic energy compared to the photon, net energy may be transferred from the electron to the photon.
- What is the energy of photon after the inverse Compton scattering? (1) In the frame K' comoving with electron, the incoming photon energy is

$$\epsilon' = \epsilon \gamma (1 - \beta \cos \theta)$$

 $\epsilon' \ll m_e c^2 = 0.5 \mathrm{MeV}$ 

17

Here,  $\theta$  is the angle between the electron velocity and the photon direction in the lab frame.

(2) In the electron rest frame, we assume the Thomson regime so that no change in the photon energy.

$$\begin{split} \epsilon_1' &= \frac{\epsilon'}{1 + \frac{\epsilon'}{mc^2} (1 - \cos \Theta')} \\ &\approx \epsilon' \left[ 1 - \frac{\epsilon'}{mc^2} (1 - \cos \Theta') \right] \quad (\text{if } \epsilon' \ll mc^2) \\ &\approx \epsilon' \quad (\text{Thomson scattering condition}) \end{split}$$
Thomson scattering condition in the rest frame:

$$\begin{array}{c} & & & & & \\ & & & \\ & & & \\ &$$

$$\mathbf{n}' = (\sin \theta' \cos \phi', \ \sin \theta' \sin \phi', \ \cos \theta')$$
$$\mathbf{n}'_1 = (\sin \theta'_1 \cos \phi'_1, \ \sin \theta'_1 \sin \phi'_1, \ \cos \theta'_1)$$
$$\cos \Theta' \equiv \mathbf{n}' \cdot \mathbf{n}'_1$$
$$= \cos \theta'_1 \cos \theta' + \sin \theta' \sin \theta'_1 \cos(\phi' - \phi'_1)$$
$$(\Theta' = \text{scattering angle in the electron rest frame})$$

(3) Going back to the lab frame, the energy of the scattered photon is

$$\epsilon_{1} = \epsilon_{1}^{\prime} \gamma \left(1 + \beta \cos \theta_{1}^{\prime}\right) \\\approx \epsilon^{\prime} \gamma \left(1 + \beta \cos \theta_{1}^{\prime}\right) \\= \epsilon \gamma^{2} \left(1 + \beta \cos \theta_{1}^{\prime}\right) \left(1 - \beta \cos \theta\right) \qquad \leftarrow \epsilon_{1}^{\prime} \approx \epsilon^{\prime} \text{ (Thomson scattering)} \\\leftarrow \epsilon^{\prime} = \epsilon \gamma \left(1 - \beta \cos \theta\right)$$

Here,  $\theta'_1$  is the scattered angle of the photon in the electron rest frame.



 $\theta_1'$ 

**→** *X* 

$$\epsilon_{1} = \epsilon \gamma^{2} \left( 1 + \beta \cos \theta_{1}' \right) \left( 1 - \beta \cos \theta \right)$$
  
=  $\epsilon \frac{1 - \beta \cos \theta}{1 - \beta \cos \theta_{1}} \qquad \leftarrow \cos \theta_{1}' = \frac{\cos \theta_{1} - \beta}{1 - \beta \cos \theta_{1}}$ (aberration

• Let's assume isotropic distribution of photons.

In the electron rest frame, most photons will be incident toward the electron, i.e.,  $\pi - \theta' \leq 1/\gamma$ .

Thomson scattering is symmetric with respect to forward and backward scattering. Therefore, in the lab frame, the scattered photon will be mostly concentrated within a narrow angle:  $\theta_1 \leq 1/\gamma$ . Assuming that  $\cos \theta_1 \approx \beta$ , we obtain  $\cos \theta_1 \approx 1 \& \beta \approx 1 \to \cos \theta_1 \approx \beta$ 

$$\epsilon_1 \approx \epsilon \frac{1 - \beta \cos \theta}{1 - \beta^2} = (1 - \beta \cos \theta) \gamma^2 \epsilon \quad \rightarrow \boxed{\langle \epsilon_1 \rangle \approx \gamma^2 \epsilon}$$
(because  $\langle \cos \theta \rangle = 0$ )

See the following slides for a precise formula.

$$\left| \left\langle \epsilon_1 \right\rangle = \gamma^2 \left( 1 + \frac{\beta^2}{3} \right) \epsilon \right|$$

Note that the radiation field is not isotropic in the electron rest frame but symmetric.





• The energies of the photon before scattering, in the electron rest frame, and after the scattering in the lab frame are in the approximate ratios

The inverse Compton scattering converts a low-energy photon to a high-energy photon by a factor of order  $\gamma^2$ 

 $\left( \epsilon : \epsilon' : \epsilon_1 \approx 1 : \gamma : \gamma^2 \right)$ 

## [Inverse Compton Power for Single Scattering]

- Assumptions:
  - (1) Isotropic distributions of photons and electrons (in the lab frame).
  - (2) The change in energy of the photon in the rest frame is negligible.

(Thomson scattering is applicable in the electron's rest frame).  $\epsilon'_1 \approx \epsilon'$ 

• Total power scattered in the electron's rest frame:

 $\frac{dE'_1}{dt'} = c\sigma_T \int \epsilon'_1 n'_{\epsilon} d\epsilon' \qquad \text{where } n'_{\epsilon} d\epsilon' \text{ is the number density of incident photons.}$ 

• Recall:  $\frac{dE_1}{dt} = \frac{dE'_1}{dt'} \quad \text{since energy and time transforms in the same way.} \\ d^3\mathbf{p} = \gamma d^3\mathbf{p'} \quad \text{transforms in the same way as energy.} \\ n_p \equiv \frac{dN}{d\mathcal{V}} \left( = \frac{d^6N}{d^3\mathbf{x}d^3\mathbf{p}} \right) \text{ is a Lorentz invariant. (density in the phase space)} \\ n_p d^3\mathbf{p} = n_\epsilon d\epsilon \\ n'_p d^3\mathbf{p'} = n'_\epsilon d\epsilon' \quad \text{because } \epsilon = pc \\ n_\epsilon d\epsilon = \gamma n'_\epsilon d\epsilon' \quad \text{Therefore, the number densities of incident photons, represented in terms of momentum and energy, transforms in the same way as energy.} \end{cases}$ 

$$\therefore \quad \frac{n_{\epsilon}d\epsilon}{\epsilon} = \frac{n_{\epsilon}'d\epsilon'}{\epsilon'}$$

Thus we have the results

 $\epsilon_1' \approx \epsilon' \ {\rm Thomson \ scattering \ assumption \ in \ the \ rest \ frame}$  $\frac{dE_1}{dt} = \frac{dE'_1}{dt'} = c\sigma_T \int \epsilon'_1 n'_{\epsilon} d\epsilon' = c\sigma_T \int \epsilon'^2 \frac{n'_{\epsilon} d\epsilon'}{\epsilon'} = c\sigma_T \int \epsilon'^2 \frac{n_{\epsilon} d\epsilon}{\epsilon}$  $= c\sigma_T \gamma^2 \int \left(1 - \beta \cos \theta\right)^2 \epsilon n_\epsilon d\epsilon$  $\leftarrow \quad \epsilon' = \epsilon \gamma (1 - \beta \cos \theta)$ 

For an isotropic distribution of photons, integrating over  $\theta$ , we have

$$\left\langle \left(1 - \beta \cos \theta\right)^2 \right\rangle = 1 + \frac{1}{3}\beta^2 \qquad \leftarrow \quad \left\langle \cos \theta \right\rangle = 0, \quad \left\langle \cos^2 \theta \right\rangle = 1/3$$

Therefore, we obtain the total power scattered in the lab frame:

 $\frac{dE_1}{dt} = c\sigma_T \gamma^2 \left(1 + \frac{1}{3}\beta^2\right) U_{\rm ph}$ 

where  $U_{\rm ph} \equiv \int \epsilon n_{\epsilon} d\epsilon$  is the initial photon energy density.

Note that the rate of decrease of the total initial photon energy is

$$\frac{dE_1^{\text{loss}}}{dt} = -c\sigma_T \int \epsilon n_\epsilon d\epsilon = -c\sigma_T U_{\text{ph}}$$

The incident power and scattered power in the lab frame can be represented in term of the rate of scattering (per unit time) and the initial and final photon energies.

$$\left|\frac{dE_1^{\text{loss}}}{dt}\right| = \epsilon \frac{dN_{\text{scatt}}}{dt}$$
$$\frac{dE_1}{dt} = \langle \epsilon_1 \rangle \frac{dN_{\text{scatt}}}{dt}$$

Then, the mean photon energy after scattering will be

$$\langle \epsilon_1 \rangle = \gamma^2 \left( 1 + \frac{1}{3} \beta^2 \right) \epsilon$$

• Thus, the net power lost by the electron, and converted into increased radiation, is

$$P_{\text{compt}} \equiv \frac{dE_1}{dt} - \left| \frac{dE_1^{\text{loss}}}{dt} \right| = c\sigma_T U_{\text{ph}} \left[ \gamma^2 \left( 1 + \frac{1}{3}\beta^2 \right) - 1 \right]$$
  
$$\therefore P_{\text{compt}} = \frac{4}{3}c\sigma_T \gamma^2 \beta^2 U_{\text{ph}} \qquad \qquad \gamma^2 - 1 = \gamma^2 \beta^2$$

• When the energy transfer in the electron rest frame is not neglected, the power is given by

$$P_{\text{compt}} = \frac{4}{3} c \sigma_T \gamma^2 \beta^2 U_{\text{ph}} \left[ 1 - \frac{63}{10} \frac{\gamma \left\langle \epsilon^2 \right\rangle}{mc^2 \left\langle \epsilon \right\rangle} \right] \quad \text{(cf. Blumenthal \& Gould, 1970)}$$

Note that the above equation allows energy to be either given or taken from the photons.

• Recall that the formula for the synchrotron power emitted by each electron is

$$P_{\text{synch}} = \frac{4}{3}\sigma_T c\gamma^2 \beta^2 U_{\text{B}}$$
  
Therefore,  $\boxed{\frac{P_{\text{synch}}}{P_{\text{compt}}} = \frac{U_{\text{B}}}{U_{\text{ph}}}}$ 

The radiation losses due to synchrotron emission and due to inverse Compton effect are in the same ratio as the magnetic field energy density and photon energy density.

• Let  $N(\gamma)d\gamma$  be the number of electrons per unit volume. Then, the total Compton power per unit volume is

$$P_{\rm tot} = \int P_{\rm compt} N(\gamma) d\gamma$$

(1) Power-law distribution of relativistic electrons ( $\beta \sim 1$ )

$$N(\gamma) = \begin{cases} C\gamma^{-p}, & \gamma_{\min} \le \gamma \le \gamma_{\max} \\ 0, & \text{otherwise} \end{cases} \longrightarrow P_{\text{tot}} = \frac{4}{3}\sigma_T c U_{\text{ph}} C (3-p)^{-1} \left(\gamma_{\min}^{3-p} - \gamma_{\max}^{3-p}\right) \end{cases}$$

(2) **Thermal distribution** of nonrelativistic electrons ( $\gamma \sim 1$ ) of number density  $n_e$ .

$$\langle \beta^2 \rangle = \langle v^2/c^2 \rangle = 3kT/mc^2 \longrightarrow P_{\text{tot}} = \left(\frac{4kT}{mc^2}\right)\sigma_T cn_e U_{\text{ph}}$$
 Note that  $\frac{4kT}{mc^2}$  is the fractional photon  $\gamma \approx 1$  energy gain because  $\frac{dE^{\text{loss}}}{dt} = -c\sigma_T U_{\text{ph}}$ .

# [Spectrum of single-scattered photons by power-law electrons]

• We will first show that a power-law photon distribution is produced from a power-law electron distribution.

The photon energy increases after a single scattering by a factor proportional to  $\gamma^2$ .

$$\langle \epsilon_1 \rangle = \gamma^2 \left( 1 + \frac{1}{3} \beta^2 \right) \epsilon = \frac{4}{3} \gamma^2 \epsilon \quad \text{as} \quad \beta \to 1$$

Therefore, the (energy) spectrum of scattered photons per unit volume per unit energy due to electrons with a power-law distribution is

 $\left(\frac{dE}{dV dt d\epsilon_1} \propto \epsilon_1^{-(p-1)/2}\right)$ 

25

The resulting spectrum has the same power-law slope as that of the synchrotron.

## [Repeated Scattering: The Compton y Parameter]

- We restrict our considerations to situations in which the Thomson limit applies:  $\gamma \epsilon \ll mc^2$
- <u>Compton *y* parameter</u>, to determine whether a photon will significantly change its energy in traversing the medium:

$$y \equiv \left(\begin{array}{c} \text{average fractional} \\ \text{energy change per} \\ \text{scattering} \end{array}\right) \times \left(\begin{array}{c} \text{mean number of} \\ \text{scatterings} \end{array}\right)$$

When  $y \gtrsim 1$ , the total photon energy and spectrum will be significantly altered; whereas for  $y \ll 1$ , the total energy is not much changed.

• Average fractional energy change per scattering (for a thermal distribution of electrons)

(a) Consider first the nonrelativistic limit (in the electron rest frame).

$$\epsilon'_1 \approx \epsilon' \left[ 1 - \frac{\epsilon'}{mc^2} \left( 1 - \cos \Theta \right) \right] \rightarrow \left\langle \frac{\Delta \epsilon'}{\epsilon'} \right\rangle \equiv \left\langle \frac{\epsilon'_1 - \epsilon'}{\epsilon'} \right\rangle = -\frac{\epsilon'}{mc^2}$$
 : angle average

In the lab frame to lowest order, this must be of the form because the mean electron energy  $\propto kT$ :

$$\left\langle \frac{\Delta \epsilon}{\epsilon} \right\rangle = -\frac{\epsilon}{mc^2} + \alpha \frac{kT}{mc^2}$$

# To calculate $\alpha$ , imagine that the photons and electrons are in complete equilibrium but interact only through scattering.

Assume that the photon density is sufficiently small that stimulated processes can be neglected. The Maxwell distribution of speeds of electrons is:

We have the averages  $\langle \epsilon \rangle$ 

$$n_{\epsilon} = K\epsilon^{2} \exp\left(-\frac{\epsilon}{kT}\right)$$
$$\langle \epsilon \rangle \equiv \int \epsilon n_{\epsilon} d\epsilon / \int n_{\epsilon} d\epsilon = 3kT$$
$$\langle \epsilon^{2} \rangle \equiv \int \epsilon^{2} n_{\epsilon} d\epsilon / \int n_{\epsilon} d\epsilon = 12(kT)^{2}$$

For this case, no net energy can be transferred from photons to electrons, so

$$\Delta \epsilon = 0 = -\frac{\left\langle \epsilon^2 \right\rangle}{mc^2} + \alpha \frac{kT}{mc^2} \left\langle \epsilon \right\rangle = \frac{3kT}{mc^2} (\alpha - 4)kT \quad \rightarrow \quad \alpha = 4$$

Thus for nonrelativisitic electrons in thermal equilibrium, the energy transfer per scattering is

$$(\Delta\epsilon)_{\rm NR} = \frac{\epsilon}{mc^2} (4kT - \epsilon)$$

Note that if the electrons have high enough temperature relative to incident photons, the photons gain energy. This is the inverse Compton scattering.

If  $\epsilon > 4kT$ , on the other hand, energy is transferred from photons to electrons.

(b) In the ultrarelativistic limit  $(\gamma \gg 1, \beta \approx 1)$ , ignoring the energy transfer in the electron rest frame,

$$\frac{P_{\rm compt}}{dE_1^{\rm loss}/dt\Big|} = \frac{4/3\sigma_T c\gamma^2 \beta^2 U_{\rm ph}}{\sigma_T cU_{\rm ph}} = \frac{4}{3}\gamma^2 \beta^2 \quad \to \quad (\Delta\epsilon)_{\rm R} \approx \frac{4}{3}\gamma^2 \epsilon$$

For a thermal distribution of ultrarelativistic electrons,

$$\langle \gamma^2 \rangle = \frac{\langle \epsilon^2 \rangle}{(mc^2)^2} = 12 \left(\frac{kT}{mc^2}\right)^2 \longrightarrow \left( (\Delta \epsilon)_{\rm R} \approx 16\epsilon \left(\frac{kT}{mc^2}\right)^2 \right)$$

$$f(\gamma) = \frac{\gamma^2 \beta}{\theta K_2(1/\theta)} e^{-\gamma/\theta}$$
  
where  $\beta = \sqrt{1 - 1/\gamma^2}$ ,  $\theta = \frac{kT}{mc^2}$ , and  
 $K_2$  is the modified Bessel function.

• Mean number of scatterings,

Recall that, for a pure scattering medium,

(mean number of scatterings) 
$$\approx \text{Max}(\tau_{\text{es}}, \tau_{\text{es}}^2)$$
 (scatterings)  $\approx \text{Max}(\tau_{\text{es}}, \tau_{\text{es}}^2)$  (where  $\tau_{\text{es}} \sim \rho \kappa_{\text{es}} R$   
 $\kappa_{\text{es}} = \frac{\sigma_T}{m_p} = 0.40 \text{ cm}^2 \text{ g}^{-1}$  for ionized hydrogen  
 $R = \text{ size of the finite medium}$ 

• <u>Compton y parameter</u>:

$$y_{\rm NR} = \frac{4kT}{mc^2} Max(\tau_{\rm es}, \ \tau_{\rm es}^2)$$

$$y_{\rm R} = 16\epsilon \left(\frac{kT}{mc^2}\right)^2 \operatorname{Max}(\tau_{\rm es}, \ \tau_{\rm es}^2)$$

- We have already shown that a power-law spectrum may be a natural consequence of a power-law distribution of electrons (single scattering case).
- We will show that a power-law photon distribution can also be produced from repeated scattering off a nonpower-law electron distribution.

Let A = the mean amplification of photon energy per scattering

 $A \equiv \frac{\epsilon_1}{\epsilon} \sim \frac{4}{3} \langle \gamma^2 \rangle$ = 16  $\left(\frac{kT}{mc^2}\right)^2$  for thermal electron distribution

initial (mean) photon energy =  $\epsilon_i$ 

(number) intensity =  $I(\epsilon_i)$  at  $\epsilon_i$ 

After k scattering, the photon energy will be  $\epsilon_k \sim \epsilon_i A^k$ .

For an optically thin scattering medium  $(\tau_{es} < 1)$ , the probability of a photon undergoing kscattering before escaping the medium is  $p_k(\tau_{es}) \sim (1 - e^{-\tau_{es}})^k \approx \tau_{es}^k$ <u>The emergent intensity at energy  $\epsilon_k$  is given by</u>  $I(\epsilon_k) \sim I(\epsilon_i) \tau_{es}^k \sim I(\epsilon_i) \tau_{es}^{\ln(\epsilon_k/\epsilon_i)/\ln A} = I(\epsilon_i) \left(\frac{\epsilon_k}{\epsilon_i}\right)^{\ln \tau_{es}/\ln A}$  $I(\epsilon_k) \sim I(\epsilon_i) \tau_{es}^k \sim I(\epsilon_i) \tau_{es}^{\ln(\epsilon_k/\epsilon_i)/\ln A} = I(\epsilon_i) \left(\frac{\epsilon_k}{\epsilon_i}\right)^{\ln \tau_{es}/\ln A}$ 

 $\therefore I(\epsilon_k) \sim I(\epsilon_i) \left(\frac{\epsilon_k}{\epsilon_i}\right)^{-\alpha} \quad \text{where} \quad \alpha \equiv \frac{-\ln \tau_{\text{es}}}{\ln A} \quad \longrightarrow \quad \text{power-law shape}$ 

• Total Compton power in the output spectrum is given by

$$P \propto \int I(\epsilon_k) d\epsilon_k = I(\epsilon_i) \epsilon_i \left[ \int x^{-\alpha} dx \right]$$

The factor in square brackets is approximately the factor by which the initial power  $I(\epsilon_i)\epsilon_i$  is amplified in energy.

Clearly, this amplification will be important if  $\alpha \ll 1$ . Therefore, energy amplification of a soft photon input spectrum is important when  $y \gtrsim 1$ :

$$\alpha = \frac{-\ln \tau_{\rm es}}{\ln A} \lesssim 1 \quad \to \quad \ln \left(\tau_{\rm es} A\right) \gtrsim 0$$
$$\rightarrow \quad y = A\tau_{\rm es} \sim 16 \left(\frac{kT}{mc^2}\right)^2 \tau_{\rm es} \gtrsim 1$$